



Laboratory Investigation Report

Patient Name : Mr. S K Sinha Centre : 3953 - Max Lab Sushant Lok 2 Sector-55

Age/Gender : 73 Y 0 M 2 D /M OP/IP No/UHID ://

 MaxID/Lab ID
 : ML03813128/3733052300115
 Collection Date/Time
 : 22/May/2023 08:35AM

 Ref Doctor
 : SELF
 Reporting Date/Time
 : 22/May/2023 08:22PM

Clinical Biochemistry

Protein/Creatinine Ratio, 24 hrs Urine

Date	22/May/2023	Unit	Bio Ref Interval
	08:35AM		
Total Volume	2200	ml/Day	1500 - 2500
Urine Protein	55.56	mg/dL	0.0-10.0
Urine Creatinine Alkaline picrate kinetic	20.11	mg/dL	
Protein/Creatinine Ratio Calculated	2.76		< 0.2

Comment 24 Hrs urine protein/creatinine ratio (PCR) is required for confirmation of ruling out significant proteinuria in positive reagent strip or positive random urine PCR. Because creatinine excretion in the urine is fairly constant throughout 24 Hrs period, measurement of PCR ratio allows correction for variation in urinary concentrations.

Proteinuria is seen in Nephrotic syndrome, diabetic nephropathy, monoclonal gammopathies and other myeloproliferative or lymphoproliferative disorders, abnormal renal tubular absorption, urinary tract malignancies, and inflammatory, degenerative and irritative conditions of the lower urinary tract.

Kindly correlate with clinical findings

*** End Of Report ***

Dr. Akash Banwari, M.D. (Path)

Dr. Jyoti Singhal, M.D. (Pathology)

Tyoti

Principal Consultant Senior Resident

Page 1 of 1

